Amendments to and Listing of the Claims:

Please cancel claims 1 and 2, without prejudice, and insert the following new claims 3-8, as set forth in the following listing of claims:

- 1. and 2. (Canceled)
- 3. (New) In a process for synthesizing a compound of formula Ib:

$$\begin{array}{c|c} R_2 & R_3 \\ \hline \\ R_4 & \end{array}$$

wherein R_1 and R_2 are each individually amino or N-alkyl substituted amino; hydroxy; alkoxy; keto; lower alkyl; or a nitrogen or oxygen protecting group;

R₃ is hydrogen; hydroxy; alkoxy; trifluoromethyl alkoxy; halo; sulfhydryl or alkylthio; R₄ is -C(O)-X;

X is hydroxy; alkoxy; or an amino acid residue;

in which process a 2-amino-5-nitro-benzonitrile starting reagent is cyclized to form 2,4-diamino-6-nitro-quinazoline, which is converted to 2,4-diamino-quinazoline, which is converted to 2,4-diamino-6-cyano-quinazoline, which is converted to 2,4-diamino-6-formyl-quinazoline;

the improvement comprising:

reacting an R₄-p-benzoic acid alkylene moiety with triethyl phosphite to form a 4-R₄-carbonyloxyalkyl-phenyl-alkyldiethylphosphite; and

reacting the 2,4-diamino-6-formyl-quinazoline with the 4-R₄-carbonyloxyalkyl-phenyl-alkyldiethylphosphite to form the compound of formula Ib.

4. (New) The process of claim 3, wherein the R_4 -p-benzoic acid alkylene moiety is a methylene moiety.

- 5. (New) The process of claim 4, wherein the R_4 -p-benzoic acid methylene moiety is 4-(bromomethyl)benzoate.
 - 6. (New) The process of claim 3, wherein X is hydroxy.
- 7. (New) The process of claim 3, wherein X is an amino acid residue of an amino acid selected from the group consisting of γ -methylene glutamic acid, glutamic acid and aspartic acid, or γ -methylene glutamate.
- 8. (New) The process of claim 3, wherein the 4-R₄-carbonyloxyalkyl-phenyl-alkyldiethylphosphite is 4-carbonyloxymethyl-phenyl-methyldiethylphosphite.
- 9. (New) The process of claim 3, further comprising hydrogenating the compound of formula Ib to form 6-(4-R₄-carbonyloxyalkylphenyl)ethanyl-2,5-diamino quinazoline, which is hydrolyzed to form 6-(4-R₄-carbonyloxyphenyl)ethanyl-2,5-diamino quinazoline,

the improvement further comprising:

reacting 6-(4-R₄-carbonyloxyphenyl)ethanyl-2,5-diamino quinazoline with diethyl γ -methylene-L-glutamate to form γ -methylene glutamate 5,8,10-trideaza aminopterin diethyl ester.

- 10. (New) The process of claim 9, wherein the improvement further comprises reacting the 6-(4-R₄-carbonyloxyphenyl)ethanyl-2,5-diamino quinazoline with the diethyl γ -methylene-L-glutamate in the presence of 1-hydroxy benzotriazole and 1-[3-dimethylamino)propyl]-3-ethyl carbodiimide hydrochloride.
- 11. (New) The process of claim 9, wherein the improvement further comprises reacting the 6-(4-R₄-carbonyloxyphenyl)ethanyl-2,5-diamino quinazoline with the diethyl γ -methylene-L-glutamate in the presence of 1-hydroxy benzotriazole, 1-[3-dimethylamino)propyl]-3-ethyl carbodiimide hydrochloride and triethylamine.